arXiv:1512.05136v2 [math.DG] 13 Apr 2016

# The Chern-Ricci flow and holomorphic bisectional curvature

Xiaokui Yang

Abstract. In this note, we show that on Hopf manifold $\mathbb{S}^{2n-1}\times\mathbb{S}^1$, the non-negativity of the holomorphic bisectional curvature is not preserved along the Chern-Ricci flow.

## 1. Introduction

The Chern-Ricci flow is an evolution equation for Hermitian metrics on complex manifolds, generalizing the Kähler-Ricci flow. Given an initial Hermitian metric $\omega_0=\sqrt{-1}(g_0)_{i\overline{j}}dz^i\wedge d\overline{z}^j$, the Chern-Ricci flow is defined as

$$\frac{\partial\omega}{\partial t}=-\mathrm{Ric}(\omega),\qquad \omega|_{t=0}=\omega_0, \tag{1.1}$$

where $\mathrm{Ric}(\omega):=-\sqrt{-1}\partial\overline{\partial}\log\det g$ is the Chern-Ricci form of $\omega$. In the case when $\omega_0$ is Kähler, namely $d\omega_0=0$, (1.1) coincides with the Kähler-Ricci flow. The Chern-Ricci flow was first introduced by Gill [3] in the setting of manifolds with vanishing first Bott-Chern classes, and many fundamental properties are established by Tosatti and Weinkove [14] on more general manifolds. A variety of further results on Chern-Ricci flow are studied in [14, 15, 16, 4, 5, 17, 18] and some of them are analogues to classical results for the Kähler-Ricci flow (e.g.[7, 2, 9, 10, 13, 11, 12]).

It is proved by Mok [8] (see [1] for Kähler threefolds and also [6]) that the non-negativity of the holomorphic bisectional curvature is preserved along the Kähler-Ricci flow. However, we show that on Hermitian manifolds, the non-negativity of the holomorphic bisectional curvature is not necessarily preserved under the Chern-Ricci flow.

**Theorem 1.1.** *Let $X=\mathbb{S}^{2n-1}\times\mathbb{S}^1$ be a diagonal Hopf manifold. Fix $T_0\geq 0$ and let*

$$\omega_0=\frac{1}{|z|^4}\sum\left((1+T_0)\delta_{ij}|z|^2-T_0\overline{z}^iz^j\right)\sqrt{-1}dz^i\wedge d\overline{z}^j.$$

*Then the Chern-Ricci flow (1.1) has maximal existence time $T_{\max}=\frac{T_0+1}{n}$.*

1. *When $t\in\left[0,\frac{T_0}{n}\right]$, $\omega(t)$ has non-negative holomorphic bisectional curvature;*
2. *However, when $t\in\left(\frac{2T_0+1}{2n},\frac{T_0+1}{n}\right)$, the holomorphic bisectional curvature of $\omega(t)$ is no longer non-negative.*

**Remark 1.2.** It is worth to point out that the same proof as in the Kähler case (following Mok) fails for the Chern-Ricci flow since the evolution of the Riemann curvature tensor under the Chern-Ricci flow involves also some terms with the torsion (and its covariant derivatives), which are not there in the Kähler-Ricci flow, where the evolution of the curvature involves only the curvature tensor itself.

**Remark 1.3.** It is also interesting to investigate sufficient conditions on Hermitian manifolds such that the non-negativity of the holomorphic bisectional curvature is preserved under the Chern-Ricci flow.

## 2. The proof of Theorem 1.1

For $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{C}^n \setminus \{0\}$ with $|\alpha_1| = \cdots = |\alpha_n| \neq 1$, let $M$ be the Hopf manifold $M = (\mathbb{C}^n \setminus \{0\})/\sim$, where

$$(z^1, \ldots, z^n) \sim \left(\alpha_1 z^1, \ldots, \alpha_n z^n\right).$$

It is easy to see that $M$ is diffeomorphic to $\mathbb{S}^{2n-1} \times \mathbb{S}^1$. Fix $T_0 > 0$ and consider the Hermitian metric

$$\omega_0 = \frac{1}{|z|^4}\left((1+T_0)\delta_{ij}|z|^2 - T_0 \overline{z}^i z^j\right)\sqrt{-1}dz^i \wedge d\overline{z}^j.$$

where $|z|^2 = \sum_{j=1}^n |z^j|^2$. It is proved in [14] that

$$\omega(t) = \omega_0 - t\mathrm{Ric}(\omega_0) \tag{2.1}$$

gives an explicit solution of the Chern-Ricci flow on $M$ with initial metric $\omega_0$. Indeed, by elementary linear algebra, we see $\det(\omega_0) = (1+T_0)^{n-1}|z|^{-2n}$ and so

$$\mathrm{Ric}(\omega_0) = n\sqrt{-1}\partial\overline{\partial}\log|z|^2 = \frac{n}{|z|^2}\left(\delta_{ij} - \frac{\overline{z}^i z^j}{|z|^2}\right)\sqrt{-1}dz^i \wedge d\overline{z}^j \geq 0.$$

For $t < \frac{T_0+1}{n}$, we have the Hermitian metrics

$$\omega(t) = \omega_0 - t\mathrm{Ric}(\omega_0) = \frac{1}{|z|^2}\left((1+T_0-nt)\delta_{ij} - (T_0-nt)\frac{\overline{z}^i z^j}{|z|^2}\right)\sqrt{-1}dz^i \wedge d\overline{z}^j. \tag{2.2}$$

Hence

$$\det(\omega(t)) = \frac{(1+T_0-nt)^{n-1}}{|z|^{2n}},$$

from which it follows that $\mathrm{Ric}(\omega(t)) = \mathrm{Ric}(\omega_0) = n\sqrt{-1}\partial\overline{\partial}\log|z|^2$. It also implies that $\omega(t)$ solves the Chern-Ricci flow on the maximal existence interval $\left[0, \frac{T_0+1}{n}\right)$.

Next, we compute the curvature tensor of the involving metric (2.2). For simplicity, we define a rescaled metric $\omega_\lambda = \sqrt{-1}h_{i\overline{j}}dz^i \wedge d\overline{z}^j$ on $M$ with

$$h_{i\overline{j}} = \frac{1}{|z|^4}\left(\delta_{ij}|z|^2 - \lambda\overline{z}^i z^j\right), \qquad \lambda < 1. \tag{2.3}$$

Note that when

$$\lambda = \frac{T_0 - nt}{1 + T_0 - nt},$$

we have

$$\omega_\lambda = \frac{\omega(t)}{1 + T_0 - nt}. \tag{2.4}$$

**Lemma 2.1.** *Let $R_{k\overline{j}i\overline{q}}$ be the curvature components of $\omega_\lambda$, then*

$$\begin{aligned} R_{k\overline{j}i\overline{q}} &= \frac{\delta_{iq}(\delta_{jk}|z|^2 - \overline{z}^k z^j)}{|z|^6} + \frac{\lambda\left(\delta_{ij}|z|^2 - \overline{z}^i z^j\right)\left(\delta_{kq}|z|^2 - \overline{z}^k z^q\right)}{|z|^8} \\ &\quad + \frac{(\lambda^2 - 2\lambda)\overline{z}^i z^q(\delta_{kj}|z|^2 - \overline{z}^k z^j)}{|z|^8}. \end{aligned}$$

*Proof.* By using elementary linear algebra, one has $\det(h_{i\overline{j}}) = (1-\lambda)|z|^{-2n}$ and so

$$\mathrm{Ric}(\omega_\lambda) = n\sqrt{-1}\partial\overline{\partial}\log|z|^2 \geq 0. \tag{2.5}$$

On the other hand, one can verify that the matrix $(h_{i\overline{j}})$ has (transpose) inverse matrix

$$h^{i\overline{j}} = |z|^2\left(\delta_{ij} + \frac{\lambda z^i \overline{z}^j}{(1-\lambda)|z|^2}\right). \tag{2.6}$$

By straightforward computation,

$$\frac{\partial h_{i\overline{j}}}{\partial z^k} = -\frac{\delta_{ij}\overline{z}^k}{|z|^4} - \frac{\lambda\delta_{jk}\overline{z}^i}{|z|^4} + \frac{2\lambda\overline{z}^i\overline{z}^k z^j}{|z|^6} = \frac{2\lambda\overline{z}^i\overline{z}^k z^j}{|z|^6} - \frac{\lambda\delta_{jk}\overline{z}^i + \delta_{ij}\overline{z}^k}{|z|^4} \tag{2.7}$$

and so

$$\begin{aligned} \Gamma^p_{ki} &= h^{p\overline{j}}\frac{\partial h_{i\overline{j}}}{\partial z^k} = |z|^2\left(\delta_{pj} + \frac{\lambda z^p\overline{z}^j}{(1-\lambda)|z|^2}\right)\left(\frac{2\lambda\overline{z}^i\overline{z}^k z^j}{|z|^6} - \frac{\lambda\delta_{jk}\overline{z}^i + \delta_{ij}\overline{z}^k}{|z|^4}\right) \\ &= \frac{2\lambda\overline{z}^i\overline{z}^k z^p}{|z|^4} - \frac{\lambda\delta_{pk}\overline{z}^i + \delta_{ip}\overline{z}^k}{|z|^2} + \frac{2\lambda^2\overline{z}^i\overline{z}^k z^p}{(1-\lambda)|z|^4} - \frac{\lambda^2\overline{z}^i\overline{z}^k z^p + \lambda\overline{z}^i\overline{z}^k z^p}{(1-\lambda)|z|^4} \\ &= \frac{\lambda\overline{z}^i\overline{z}^k z^p}{|z|^4} - \frac{\lambda\delta_{pk}\overline{z}^i + \delta_{ip}\overline{z}^k}{|z|^2}. \end{aligned}$$

The Chern curvature tensor of $\omega_\lambda$ is

$$\begin{aligned} R^p_{k\overline{j}i} &= -\frac{\partial\Gamma^p_{ki}}{\partial\overline{z}^j} \\ &= -\frac{\lambda\delta_{ij}\overline{z}^k z^p + \lambda\delta_{kj}\overline{z}^i z^p}{|z|^4} + \frac{2\lambda\overline{z}^i\overline{z}^k z^p z^j}{|z|^6} + \frac{\lambda\delta_{pk}\delta_{ij} + \delta_{ip}\delta_{kj}}{|z|^2} - \frac{\lambda\delta_{pk}\overline{z}^i z^j + \delta_{ip}\overline{z}^k z^j}{|z|^4} \\ &= \frac{\lambda\delta_{pk}\delta_{ij} + \delta_{ip}\delta_{kj}}{|z|^2} + \frac{2\lambda\overline{z}^i\overline{z}^k z^p z^j}{|z|^6} - \frac{\lambda\left(\delta_{ij}\overline{z}^k z^p + \delta_{kj}\overline{z}^i z^p + \delta_{pk}\overline{z}^i z^j\right) + \delta_{ip}\overline{z}^k z^j}{|z|^4}. \end{aligned}$$

Hence

$$\begin{aligned}
& R_{k\bar{j}i\bar{q}} = h_{p\bar{q}} R^p_{k\bar{j}i} \\
= \; & \frac{\delta_{pq}|z|^2}{|z|^4}\left[\frac{\lambda\delta_{pk}\delta_{ij}+\delta_{ip}\delta_{kj}}{|z|^2}+\frac{2\lambda\overline{z}^i\overline{z}^k z^p z^j}{|z|^6}-\frac{\lambda\left(\delta_{ij}\overline{z}^k z^p+\delta_{kj}\overline{z}^i z^p+\delta_{pk}\overline{z}^i z^j\right)+\delta_{ip}\overline{z}^k z^j}{|z|^4}\right] \\
& -\frac{\lambda\overline{z}^p z^q}{|z|^4}\left[\frac{\lambda\delta_{pk}\delta_{ij}+\delta_{ip}\delta_{kj}}{|z|^2}+\frac{2\lambda\overline{z}^i\overline{z}^k z^p z^j}{|z|^6}-\frac{\lambda\left(\delta_{ij}\overline{z}^k z^p+\delta_{kj}\overline{z}^i z^p+\delta_{pk}\overline{z}^i z^j\right)+\delta_{ip}\overline{z}^k z^j}{|z|^4}\right] \\
= \; & \frac{\lambda\delta_{qk}\delta_{ij}+\delta_{iq}\delta_{jk}}{|z|^4}+\frac{2\lambda\overline{z}^i\overline{z}^k z^j z^q}{|z|^8}-\frac{\lambda\left(\delta_{ij}\overline{z}^k z^q+\delta_{kj}\overline{z}^i z^q+\delta_{kq}\overline{z}^i z^j\right)+\delta_{iq}\overline{z}^k z^j}{|z|^6} \\
& -\frac{\lambda^2\delta_{ij}\overline{z}^k z^q+\lambda\delta_{kj}\overline{z}^i z^q}{|z|^6}-\frac{2\lambda^2\overline{z}^i\overline{z}^k z^j z^q}{|z|^8} \\
& +\frac{\lambda^2\left(\delta_{ij}\overline{z}^k z^q|z|^2+\delta_{kj}\overline{z}^i z^q|z|^2+\overline{z}^i\overline{z}^k z^j z^q\right)+\lambda\overline{z}^i\overline{z}^k z^j z^q}{|z|^8} \\
= \; & \frac{\lambda\delta_{qk}\delta_{ij}+\delta_{iq}\delta_{jk}}{|z|^4}+\frac{(3\lambda-\lambda^2)\overline{z}^i\overline{z}^k z^j z^q}{|z|^8}-\frac{\lambda\delta_{qk}\overline{z}^i z^j}{|z|^6}-\frac{\lambda\delta_{ij}\overline{z}^k z^q}{|z|^6} \\
& +\frac{(\lambda^2-2\lambda)\delta_{kj}\overline{z}^i z^q}{|z|^6}+\frac{\delta_{iq}\overline{z}^k z^j}{|z|^6} \\
= \; & \frac{\delta_{iq}(\delta_{jk}|z|^2-\overline{z}^k z^j)}{|z|^6}+\frac{\lambda\delta_{ij}\left(\delta_{kq}|z|^2-\overline{z}^k z^q\right)}{|z|^6}+\frac{(\lambda^2-2\lambda)\overline{z}^i z^q(\delta_{kj}|z|^2-\overline{z}^k z^j)}{|z|^8} \\
& +\frac{\lambda\overline{z}^i z^j\left(\overline{z}^k z^q-\delta_{kq}|z|^2\right)}{|z|^8} \\
= \; & \frac{\delta_{iq}(\delta_{jk}|z|^2-\overline{z}^k z^j)}{|z|^6}+\frac{\lambda\left(\delta_{ij}|z|^2-\overline{z}^i z^j\right)\left(\delta_{kq}|z|^2-\overline{z}^k z^q\right)}{|z|^8}+\frac{(\lambda^2-2\lambda)\overline{z}^i z^q(\delta_{kj}|z|^2-\overline{z}^k z^j)}{|z|^8}.
\end{aligned}$$

□

**Lemma 2.2.** *For any $\lambda \in [0,1)$, $\omega_\lambda$ has non-negative holomorphic bisectional curvature.*

*Proof.* For any $\xi = (\xi^1, \cdots, \xi^n)$ and $\eta = (\eta^1, \cdots, \eta^n)$, by Lemma 2.1 we have

$$\begin{aligned}
R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\eta^i\overline{\eta}^q \;=\; & \frac{|\eta|^2(|z|^2|\xi|^2-|\overline{z}\cdot\xi|^2)}{|z|^6}+\frac{\lambda\left|\left(\delta_{ij}|z|^2-\overline{z}^i z^j\right)\eta^i\overline{\xi}^j\right|^2}{|z|^8} \\
& +\frac{(\lambda^2-2\lambda)|\overline{z}\cdot\eta|^2(|z|^2|\xi|^2-|z\cdot\overline{\xi}|^2)}{|z|^8}.
\end{aligned}$$

Since $|z|^2|\eta|^2 \geq |\overline{z}\cdot\eta|^2$, we obtain

$$R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\eta^i\overline{\eta}^q \geq \frac{\lambda\left|\left(\delta_{ij}|z|^2-\overline{z}^i z^j\right)\eta^i\overline{\xi}^j\right|^2}{|z|^8}+\frac{(\lambda^2-2\lambda+1)|\overline{z}\cdot\eta|^2(|z|^2|\xi|^2-|z\cdot\overline{\xi}|^2)}{|z|^8}.$$

The right hand side is non-negative when $\lambda \geq 0$. □

**Corollary 2.3.** *The initial metric $\omega_0$ has non-negative holomorphic bisectional curvature.*

*Proof.* When $t=0$, or equivalently $\lambda=\frac{T_0}{1+T_0}$, we know $\omega_\lambda=\frac{\omega_0}{1+T_0}$. Since $\lambda=\frac{T_0}{1+T_0}\in[0,1)$, by Lemma 2.2, $\omega_0$ has non-negative holomorphic bisectional curvature. □

**Lemma 2.4.** *When $\lambda<-1$, the holomorphic sectional curvature of the metric $\omega_\lambda$ is no longer non-negative. In particular, the holomorphic bisectional curvature of the metric $\omega_\lambda$ is no longer non-negative.*

*Proof.* For any $\xi=(\xi^1,\cdots,\xi^n)$, we have

$$\begin{aligned}R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\xi^i\overline{\xi}^q &= \frac{|\xi|^2(|z|^2|\xi|^2-|\overline{z}\cdot\xi|^2)}{|z|^6}+\frac{\lambda(|z|^2|\xi|^2-|\overline{z}\cdot\xi|^2)^2}{|z|^8}\\ &\quad+\frac{(\lambda^2-2\lambda)|\overline{z}\cdot\xi|^2(|z|^2|\xi|^2-|\overline{z}\cdot\xi|^2)}{|z|^8}\\ &= \frac{(3\lambda-\lambda^2)|\overline{z}\cdot\xi|^4+(\lambda+1)(|z|^2|\xi|^2)^2+(\lambda^2-4\lambda-1)|\overline{z}\cdot\xi|^2|z|^2\cdot|\xi|^2}{|z|^8}.\end{aligned}$$

Let $a=|\overline{z}\cdot\xi|^2$ and $b=|z|^2|\xi|^2$, then

$$\begin{aligned}R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\xi^i\overline{\xi}^q &= \frac{(3\lambda-\lambda^2)a^2+(\lambda^2-4\lambda-1)ab+(\lambda+1)b^2}{|z|^8}\\ &= \frac{(b-a)a(\lambda-1)^2+(b-a)^2(\lambda+1)}{|z|^8}.\end{aligned}$$

It is easy to see that, $b\geq a\geq 0$ and so for any $-1\leq\lambda<1$

$$R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\xi^i\overline{\xi}^q\geq 0.$$

However, when $\lambda<-1$, $R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\xi^i\overline{\xi}^q$ is no longer nonnegative. Indeed, for any given $z=(z^1,\cdots,z^n)$, we choose a nonzero vector $\xi=(\xi^1,\cdots,\xi^n)$ such that $\overline{z}\cdot\xi=0$, i.e. $\sum\overline{z}^i\cdot\xi^i=0$. In this case, we have $a=|\overline{z}\cdot\xi|=0$, but $b=|z|^2|\xi|^2>0$. Moreover,

$$R_{k\bar{j}i\bar{q}}\xi^k\overline{\xi}^j\xi^i\overline{\xi}^q=\frac{b^2(\lambda+1)}{|z|^8}<0$$

since $\lambda<-1$. □

*The proof of Theorem 1.1.* By (2.4), we see when $\lambda=\frac{T_0-nt}{1+T_0-nt}$, $\omega_\lambda=\frac{\omega(t)}{1+T_0-nt}$. Hence,

(1) by Lemma 2.2, when $\lambda\in[0,1)$ or equivalently, $0\leq t\leq\frac{T_0}{n}$, $\omega(t)$ has non-negative holomorphic bisectional curvature;
(2) by Lemma 2.4, when $\lambda<-1$, or equivalently, $\frac{2T_0+1}{2n}<t<\frac{T_0+1}{n}$, the holomorphic bisectional curvature of $\omega(t)$ is no longer non-negative. □

**Acknowledgements.** This work was partially supported by China's Recruitment Program of Global Experts, as well as National Center for Mathematics and Interdisciplinary Sciences, Chinese Academy of Sciences. The author wishes to thank Professor Ben Weinkove and Professor Valentino Tosatti for their invaluable support and guidance. The author is grateful to the referees for very careful reading and many helpful suggestions.

## References

[1] Bando, S. On the classification of three-dimensional compact Kähler manifolds of nonnegative bisectional curvature. J. Differential Geom. **19** (1984), no. 2, 283–297.

[2] Cao, H.-D. Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds, Invent. Math. **81** (1985), no. 2, 359–372.

[3] Gill, M. Convergence of the parabolic complex Monge-Ampère equation on compact Hermitian manifolds, Comm. Anal. Geom. **19** (2011), no. 2, 277–303.

[4] Gill, M. The Chern-Ricci flow on smooth minimal models of general type. arXiv:1307.0066

[5] Gill, M.; Smith, D. The behavior of Chern scalar curvature under Chern-Ricci flow. arXiv:1311.6534.

[6] Gu, H.-L. A new proof of Mok's generalized Frankel conjecture theorem. Proc. Amer. Math. Soc. **137** (2009), no. 3, 1063–1068.

[7] Hamilton, R.S. Three-manifolds with positive Ricci curvature, J. Differential Geom. **17** (1982), no. 2, 255–306.

[8] Mok, N. The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature. J. Differential Geom. **27** (1988), no. 2, 179–214.

[9] Song, J.; Tian, G. The Kähler-Ricci flow on surfaces of positive Kodaira dimension, Invent. Math. **170** (2007), no. 3, 609–653.

[10] Song, J.; Tian, G. Canonical measures and Kähler-Ricci flow, J. Amer. Math. Soc. **25** (2012), no. 2, 303–353.

[11] Song, J.; Weinkove, B. Contracting exceptional divisors by the Kähler-Ricci flow, Duke Math. J. **162** (2013), no. 2, 367–415.

[12] Song, J.; Weinkove, B. Lecture notes on the Kähler-Ricci flow, arXiv:1212.3653.

[13] Tian, G.; Zhang, Z. On the Kähler-Ricci flow on projective manifolds of general type, Chinese Ann. Math. Ser. B **27** (2006), no. 2, 179–192.

[14] Tosatti, V.; Weinkove, B. On the evolution of a Hermitian metric by its Chern-Ricci form, J. Differential Geom. **99** (2015), no. 1, 125–163.

[15] Tosatti, V.; Weinkove, B. The Chern-Ricci flow on complex surfaces, Compos. Math. **149** (2013), no. 12, 2101–2138.

[16] Tosatti, V.; Weinkove, B.; Yang, X.-K. Collapsing of the Chern-Ricci flow on elliptic surfaces. Math. Ann. **362** (2015), no.3, 1223–1271.

[17] Fang, S.-W; Tosatti, V.; Weinkove, B.; Zheng, T. Inoue surfaces and the Chern-Ricci flow. arXiv:1501.07578.

[18] Zheng, T. *The Chern-Ricci flow on Oeljeklaus-Toma manifolds.* arXiv:1505.07299.

Current address of Xiaokui Yang: Morningside Center of Mathematics, Institute of Mathematics, Hua Loo-Keng Key Laboratory of Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, 100190, China.

*E-mail address*: xkyang@amss.ac.cn